



National
Library
of Medicine



My NC
[Sign In] [Regis]

All Databases

PubMed

Nucleotide

Protein

Genome

Structure

OMIM

PMC

Journals

Book

Search for

Limits Preview/Index History Clipboard Details

Display Show Sort by Send to

About Entrez

Text Version

Entrez PubMed

Overview

Help | FAQ

Tutorial

New/Noteworthy

E-Utilities

PubMed Services

Journals Database

MeSH Database

Single Citation Matcher

Batch Citation Matcher

Clinical Queries

Special Queries

LinkOut

My NCBI (Cubby)

Related Resources

Order Documents

NLM Mobile

NLM Catalog

NLM Gateway

TOXNET

Consumer Health

Clinical Alerts

ClinicalTrials.gov

PubMed Central

☐ 1: Dev Biol Stand. 1998;92:3-11.

[Related Articles, Links](#)

The mode of action of immunological adjuvants.

Allison AC.

Dawa Corporation, Belmont, CA, USA.

Adjuvants augment immune responses to antigens and influence the balance between cell-mediated and humoral responses, as well as the isotypes of antibodies formed. New adjuvant formulations include antigen-carrying vehicles and small molecules with immunomodulating activity. Widely used two-phase vehicles comprise liposomes and microfluidized squalene or squalene emulsions. These are believed to target antigens to antigen-presenting cells, including dendritic cells (DC), follicular dendritic cells (FDC) and B-lymphocytes. Activation of complement generates C3d, which binds CR2 (CD21) on FDC and B-lymphocytes, thereby stimulating the proliferation of the latter and the generation of B-memory. Targeting of antigens to DC may favour cell-mediated immunity. Immunomodulating agents induce the production of cytokine cascades. In a primary cascade at injection sites TNF-alpha, GM-CSF and IL-1 are produced. TNF-alpha promotes migration of DC to lymphoid tissues, while GM-CSF and IL-1 accelerate the maturation of DC into efficient antigen-presenting cells for T-lymphocytes. In a secondary cytokine cascade in draining lymph nodes, DC produce IL-12, which induces Th1 responses with the production of IFN-gamma. The cytokines elicit cell-mediated immune responses and the formation of antibodies of protective isotypes, such as IgG2a in the mouse and IgG1 in humans. Antibodies of these isotypes activate complement and collaborate with antibody-dependent effector cells in protective immune responses.

Publication Types:

- Review
- Review, Tutorial

PMID: 9554254 [PubMed - indexed for MEDLINE]

Display ☐ Show ☐ Sort by ☐ Send to ☐

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)

Aug 23 2005 04:56:19